

Phytoplankton: The Life Jacket of the Planet Earth

Abhijit Mitra

Department of Marine Science, University of Calcutta, India

Article Info

***Corresponding author:**

Abhijit Mitra

Department of Marine Science
University of Calcutta
35, B.C. Road, Kolkata 700 019
India
E-mail: abhijit_mitra@hotmail.com

Received: November 16, 2023

Accepted: December 04, 2023

Published: December 20, 2023

Citation: Mitra A. Phytoplankton: The Life Jacket of the Planet Earth. *Madridge J Agric Environ Sci.* 2023; 5(1): 68-70.
doi: 10.18689/mjaes-1000113

Copyright: © 2023 The Author(s). This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Published by Madridge Publishers

The pelagic environment of the ocean supports two basic types of marine organisms. One type comprises the **plankton**, or those organisms whose powers of locomotion are such that they are incapable of making their way against the current and thus are passively transported by currents in the aquatic system and the other type includes the **nekton** (free swimmers), which are free-floating animals that, in contrast to plankton, are strong enough to swim against currents and are therefore independent of water movements. The category of nekton includes fish, squid and marine mammals.

The word plankton has come from the Greek word "*planktos*", meaning that which is passively drifting or wandering. Depending upon whether a planktonic organism is a plant or animal, a distinction is made between **phytoplankton** and **zooplankton**. Although many planktonic species are of microscopic dimensions, the term is not synonymous with small size as some of the zooplankton includes jellyfish of several meters in diameter. It is not necessary that all plankton is completely passive, most of them are capable of swimming too.

Phytoplankton are free floating tiny floral components that are widely distributed in the marine and estuarine environments. Like land plants, these tiny producers require sunlight, nutrients or fertilizers, carbon dioxide gas and water for growth. The cells of these organisms contain the pigment **chlorophyll** that traps the solar energy for use in **photosynthesis**. The photosynthetic process uses the solar radiation to convert carbon dioxide and water into sugars or high energy organic compounds from which the cell forms new materials. The synthesis of organic material by photosynthesis is termed **primary production**. Since phytoplankton are the dominant producers in the ocean, their role in the marine food chain is of paramount importance. Approximately, 4000 species of marine phytoplankton have been described and new species are continually being added to this total. Phytoplankton exhibit remarkable adaptations to remain in floating condition in the seawater. In fact, all marine phytoplankton tend to stay in the photic zone to utilize the solar radiation for performing the process of photosynthesis. In order to retard the process of sinking, this group of organisms adopts various mechanisms. These include their small size and general morphology, as the ratio of cell surface area to volume determines frictional drag in the water. Colony or chain formation also increases surface area and slows sinking. Most species carry out ionic regulation, in which the internal concentration of ions is reduced relative to their concentration in seawater. Diatoms also produce and store oil and this metabolic by-product further reduces cell density. In experimental conditions, living cells tend to sink at rates ranging from 0 to 30 m day⁻¹, but dead cells may sink more than twice as fast. In nature, turbulence of surface waters is also an important factor in maintaining phytoplankton near the surface where they receive abundant sunlight.

Phytoplankton present in the marine and estuarine environments use carbon dioxide for photosynthesis and hence play an important role in maintaining the carbon dioxide budget of the atmosphere. The larger the world's phytoplankton population, the more

carbon dioxide gets pulled from the atmosphere. This lowers the average temperature of the atmosphere due to lower volumes of this greenhouse gas. Scientists have found that a given population of phytoplankton can double its numbers in the order of once per day. In other words, phytoplankton responds very rapidly to changes in their environment.

Phytoplankton sometimes may cause adverse impact on the marine and estuarine environment. During excessive bloom of phytoplankton, the light energy is intercepted, which could otherwise reach fixed plants like eel grass (*Zostera* spp.) and kelp. Furthermore, when the phytoplankton eventually die back and break down, an excessive amount of oxygen is required to fuel this process, and hence areas may become deprived of oxygen. Excessive nutrients, and/ or changes in their relative concentrations, may be one factor in a chain of events leading to changes in the species composition of the phytoplankton communities. Increased occurrence of toxic algal blooms may accelerate toxin production. Toxic phytoplankton, when consumed by shellfish or other species can affect the marine food chain, including poisoning of seabirds, mammals and even humans. It has been established that phytoplankton naturally contains DMS (dimethyl sulphide), which is released from dead phytoplankton into the atmosphere. This compound can transform into sulphuric acid, which eventually may contribute to acid rain (<http://oceanlink.island.net/ask/pollution.html>).

Nearly all marine plants, whether unicellular or multicellular, even those attached to substrata (sessile) or free floating, pass some part of their life cycle in floating condition as phytoplankton. However, those organisms which always remain planktonic throughout the life cycle are 1) diatoms, 2) dinoflagellates, 3) coccolithophores, 4) selective species of blue-green algae and 5) some species of green algae.

Diatoms

These floating plants are all microscopic in size and are characterised by the presence of shell or frustule. The shell or frustule is composed of translucent silica. The cell wall of diatom has two parts resembling a pillbox bottom and lid. The lid is called the **epitheca** and the bottom is known as **hypotheca**. These shells have great importance from the geological point of view and constitute the diatomaceous crust. The diatoms exhibit remarkable varieties and forms and many species possess beautifully sculptured shells.

Depending on the nature of valves and pattern of ornamentation in the valve surface, the diatoms are grouped into **centric** and **pennate** diatoms. The major differences between these two groups are given in table 1.

Table 1. Differences between centric and pennate diatoms

| Point | Centric diatom | Pennate diatom |
|---------------|--|--|
| Cell shape | Discoïd, solenoid or cylindrical. | Elongated and fusiform, oval, sigmoid, or roughly circular. |
| Ornamentation | Radial in nature i.e., the arrangement of markings is radiating from the centre. | Bilateral in nature i.e., the arrangement of the markings is on either side of the apical (main) axis. |

Dinoflagellates

These are important producers of the marine environment and rank second in importance in the economy of the sea. Typically, these are unicellular, some are naked while others are armoured with plates of cellulose. The dinoflagellates possess two flagella for locomotion. Several of them are luminescent and produce light. *Skeletonema costatum* and *Coscinodiscus eccentricus* are common dinoflagellates in marine and estuarine system.

Coccolithophores

These are among the smallest category of phytoplankton having a size range between 5 to 20 microns. Some coccolithophores have flagella while others are devoid of them. Their soft bodies are shielded by tiny, calcified circular plates or shields of various designs. These are normally found in the open sea, but their profuse occurrence has been recorded in coastal waters. They form important diet components of filter feeding animals. Examples are *Isochrysis galbana* and *Coccolithus* sp.

Blue-Green Algae

These include both unicellular and multicellular organisms. The blue colour in them is due to the presence of a pigment known as phycocyanin. Of the various organisms belonging to this category, the most important is *Trichodesmium erythraeum* because in certain seasons of the year its biomass increases greatly resulting in the formation of clumps.

Green Algae

Microscopic green algae present in the planktonic community largely occur in coastal waters. The green colour in them is due to the presence of chloroplasts. They are widely distributed in the warmer (tropical) seas and only few species are found in the Arctic and Antarctic oceans. *Chlorella marina* and *Chlorella salina* are common green algae in estuarine waters.

Phytoplankton: The Service Provider in Context to Climate Change

If trees are the lungs of the land, phytoplankton are the lungs of the ocean, producing about half of the world's oxygen. While trees have full leaf canopies to collect carbon from the atmosphere and turn it into oxygen, phytoplankton are microscopic and rely on sheer numbers to help transport 9 trillion kilograms of carbon (equivalent to the weight of over 5 million blue whales) from the atmosphere to the depths of the ocean. Approximately 25,000 species of phytoplankton (documented till date) worldwide work in harmony to photosynthesize and create food that is the basis for almost all marine food webs.

"Without phytoplankton, the world would be in serious trouble. As humans continue to change the ocean, it is *vital* to understand how phytoplankton will be affected. So, to understand how phytoplankton will respond to a changing

ocean, researchers consolidated large number of scientific studies on phytoplankton and changes in nutrients with ocean acidification and warming. The researchers found that with a more acidic ocean, phytoplankton had a dramatic change in how they used nutrients like becoming more efficient in their use of nitrogen when nitrogen was limited. When the phytoplankton were put under even more stress in nutrient limited conditions, the change was even greater. Essentially, phytoplankton changed the way their cells were functioning due to stress, just like how many people tend to eat more when suffer from stress or depression.

The ocean is warming just like the atmosphere. Each phytoplankton species is used to a different temperature – those in the arctic like it cold, while those in the tropics prefer tank top weather. However, researchers found that most phytoplankton grew better with warmer ocean temperatures.

Not only does this mean that phytoplankton will grow in heat, but it also means that they will be able to reproduce more quickly, creating even more phytoplankton. Scientists think that warmer oceans would make it easier for phytoplankton cells to function and would decrease the amount of phosphorous each plankton needs. Arctic and temperate regions are expected to feel the largest impacts of ocean heating in the future, leading to the largest change in phytoplankton dynamics in these areas.

These creatures may be microscopic in size, but, when taken together, this tiny producer community has an enormous regulatory influence on the planet's atmosphere, capturing as much as 40% of all CO₂ produced, which is roughly about four times the amount captured by the Amazon rainforest.