

2nd International Conference on

NEUROLOGY AND NEUROSURGERY

December 11, 2020 | Virtual Conference

Spatiotemporal Dynamics of Activity Patterns in Neurological Disorders

Radu Mutihac

University of Bucharest, Romania

Connectivity studies using resting-state functional magnetic resonance imaging (rsfMRI), diffusion tensor imaging (DTI) and more recently, diffusion spectroscopic imaging (DSI) have enhanced our knowledge on the organization of large-scale structural and functional brain networks, which consist of spatially distributed, but functionally linked regions that continuously share information with each other. Brain's energy is largely consumed at rest during spontaneous neuronal activity (~20%), while task-related increases in metabolism energy are minor (<5%). Spontaneous ultralow-frequency fluctuations in BOLD-based rsfMRI signals (<0.08Hz) at the level of large-scale neural systems are not noise, but orderly and organized in a series of functional networks that permanently maintain a high level of temporal coherence among brain areas that are structurally segregated and functionally linked in resting state networks (RSNs). Some RSNs are functionally organized as dynamically competing systems both at rest and during tasks. The default mode network (DMN), the most important RSN, is even more active during rest and involved in realization of tasks like memory retrieval, emotional process and social cognition. Cortical connectivity at rest is reportedly altered in several neurological and psychiatric disorders. Most recently, human brain function has been imaged in fMRI and thereby accessing both sides of the mind-brain interface (subjective experience and objective observations) have simultaneously been performed. As such functional neuroimaging moves on to new potential applications like reading the brain states, brain-computer interfaces, lie detection and so forth. The presentation aims to review and evaluate the most current approaches and findings on early detection and classification of cognitive impairments and dementia, particularly among syndromes with relatively similar behavioral effects, on the basis of alterations in brain connectivity at rest explored by fused rsfMRI, DTI and DSI.