



# International Conference on Traditional Medicine and Ethnobotany

September 23-24, 2019 Kuala Lumpur, Malaysia

## The Relationship between the Spasticity-Related Pain and the Quality of Life about the Cerebral Palsy in China

Zhenhuan Liu<sup>1\*</sup>, Yong Zhao<sup>1</sup>, Wenjie Fu<sup>1</sup>, Xuguang Qian<sup>1</sup>, Chuntao Zhang<sup>1</sup>, Shangsheng Dong<sup>2</sup>, Mona L. Martin<sup>3</sup>, Donald M. Bushnell<sup>3</sup>, Thorin L. Geister<sup>4</sup>, Yuqiong Zhang<sup>5</sup> and Jie Yang<sup>6</sup>

<sup>1</sup>Guangzhou University of Traditional Chinese Medicine, China

<sup>2</sup>Jiangmen Maternity and Child Health Care, China

<sup>3</sup>Health Research Associates, USA

<sup>4</sup>Merz Pharmaceuticals GmbH Eckenheimer, Germany

<sup>5</sup>Rehabilitation Division MCH Hospital of Dongguan, China

<sup>6</sup>XiangYaBoAi Rehabilitation Hospitals, China

**Purpose:** Children and adolescents with cerebral palsy suffered from spastic pain that affected their quality of life. The purpose of this study was to examine the relationship between the spasticity-related pain and the quality of life about the children with cerebral palsy with spasticity-related pain in China.

**Methods:** This was a prospective, observational, one-arm multicenter study. 137 cerebral palsy children with SRP and their parents/caregivers were recruited from three sites in China. The children and their parents/caregivers completed the Questionnaire on pain caused by Spasticity and PedsQL™ 4 scale.

**Results:** 137 children with their parents/caregivers were eligible for participation in the study and all completed the relevant investigation. According to the result of the children-report PedsQL, the total scale scores in the V2/V3 about the different age groups (5-7 years old, 8-12 years old, 13-16 years old) had no statistical significance ( $P > 0.05$ ). After 3 weeks, the change compared to first time survey result was not significant ( $p > 0.05$ ). The total scores and each domain of PedsQL in the CP with SRP was lower than the normal group ( $P < 0.05$ ). On the whole, the associations between the child and parent proxy-report QPS total scores and each domain scores of PedsQL were moderate to high. The PedsQL total scores relationship to the QPS item score was moderate.

**Conclusion:** The quality of life in the CP patients with SRP was worse than normal people and it was a steady long-term stage. The correlation between the spasticity-related pain and the quality of life was moderate to high and negative and the result from the parent proxy-report was more evident.